

VISUAL WIDE BINARIES AND THE STRUCTURE OF PLANETARY NEBULAE

Noam Soker

Department of Physics, University of Haifa at Oranim

Oranim, Tivon 36006, ISRAEL

soker@physics.technion.ac.il

ABSTRACT

In a recent work Ciardullo *et al.* (1999) list 19 planetary nebulae surveyed by the Hubble Space Telescope for the presence of resolved visual binary companions of their central stars. For ten planetary nebulae they argue for probable physical association of the resolved stars with the central stars, while for nine the association is less likely. Such stellar companions, at orbital separations of hundreds to thousands of astronomical units, will cause the structures of these planetary nebulae to possess a non-axisymmetrical signatures. By using images from the literature of these 19 planetary nebulae, I demonstrate that the structures of the planetary nebulae are compatible in most cases with Ciardullo *et al.*'s arguments for an association, or not, of the resolved stars with the planetary nebulae central stars. This shows that the departure, or not, of a planetary nebula from having pure axisymmetrical structure can be used to strengthen an argument for an association, or a non-association, of a putative wide companion with the stellar progenitor of the nebula.

Subject heading: Planetary nebulae:general – binaries: visual – stars: AGB and post-AGB

1. INTRODUCTION

In a recent paper Ciardullo *et al.* (1999) resolved the companions (or possible companions) of 19 planetary nebulae (PNs) nuclei. By assuming that the resolved companions are main sequence stars, they calculated the distances to these PNs. The projected orbital separations of the binary nuclei are in the range of a few hundred to several thousand astronomical units. At such large orbital separations the companions will not influence the mass loss from the PN progenitors. However, the companions can influence the structures of the descendant PNs. For orbital periods of $P_{\text{orb}} \lesssim 10\tau_{PN}$, where τ_{PN} is the formation time for the nebula, the orbital motion may cause a departure from axisymmetrical structure (Soker 1994), while for larger separations the companion may blow a small bubble inside the nebula (Soker 1996). Such a bubble, formed by a very wide companion, might be the “vertical bridge” observed by Corradi *et al.* (1999) near a star located in Wray 17-1.

The list of binary companions presented by Ciardullo *et al.* allows a qualitative comparison between the properties of the binary nuclei and the observed structures of these PNs. This is the aim of this paper: to search for a consistency between the claimed binary nuclei and the PNs morphologies, thereby to support the arguments of Ciardullo *et al.* (1999) for an association, or not, between the observed companions and the central stars of the PNs. As an example, let us examine the Stingray nebula, for which a visual companion was resolved by the Hubble space telescope near its central star (Bobrowsky *et al.* 1998). The projected orbital separation is ~ 2200 AU, while the outer nebular radius is ~ 8000 AU. This radius implies a nebular expansion age of $\tau_{PN} \sim 4000$ years for an expansion velocity of 10 km s^{-1} . A companion at such a large separation cannot be responsible for the axisymmetrical structure of this PN. Its orbital separation, which implies an orbital period of $\sim 10^5$ yrs, may manifest itself in a departure from axisymmetry. During the time of $\tau_{PN} \simeq 4000$, the mass-losing star would have moved a distance of $\sim 0.5(4000/10^5) \times 2\pi \times 2200 \text{ AU} \simeq 275 \text{ AU}$ in its orbital motion (assuming the two stars have equal masses). This is $\sim 3\%$ of the nebular radius, and it implies a clear deviation from axisymmetry. This PN, indeed, has a large degree of deviation from axisymmetry, part of

which may be attributed to the presence of this companion. However, this PN has a complicated structure, including a dense ring which is inclined to the long axis of the nebula. The overall degree of deviation from axisymmetry seems to me to be too large to be solely explained by the visual companion, and I hypothesize that a closer companion in an eccentric orbit may exist. The visually resolved companion’s wind may also have shaped the nebula in its immediate vicinity. The Stingray nebula deserves further observations.

The paper is organized as follows. In §2 I find the masses of the main sequence companions, and calculate the orbital periods and velocities. I then mark each PN according to whether its structure is compatible or not with the properties of its central binary system, and in §3 I elaborate on the individual PNs. In §4 I compare the sample studied in §3 to the sample of PNs for which Ciardullo *et al.* found no companions. A short summary is in §5.

2. PROPERTIES OF BINARY NUCLEI

In Table 1 the same 19 PNs that are given by Ciardullo *et al.* (1999) in their table 7 are listed. The columns in Table 1 are: (1) The common PN name. (2) The PN G designation (galactic coordinates). (3) The secondary mass in $M_{\odot}$. It is calculated from the magnitude M_V , which is found from the colors (V-I) and reddening $[E(B-V)]$ as given by Ciardullo *et al.* Only for K 1-27 was the companion’s mass taken to be $0.6M_{\odot}$ since Ciardullo *et al.* argue that the companion is a WD rather than a main sequence star. (4) The projected orbital separation in AU. The values are taken from Ciardullo *et al.*, where more details can be found. (5) The orbital period, in years, calculated by taking the progenitor of the PN (now the WD) to be of mass $0.8M_{\odot}$. The real orbital separation is likely to be larger than the projected orbital separation. An effect in the opposite direction is the mass-loss process during the AGB phase. The orbital separation increases during this phase, and hence it was smaller at the beginning of the process. The projection effect is likely to dominate in most cases, hence the orbital periods are likely to be longer than those given in the table. (6) The orbital velocity of the primary around the binary center of mass, taking the projected orbital separation and assuming a circular orbit and a mass

of $0.8M_{\odot}$ for the WD progenitor. (7) The nebular diameter as given by Ciardullo *et al.* (1999), in parsecs. (8) The classification of Ciardullo *et al.* (1999; Table 7): P–Probable binary; S–Possible binary; D–Doubtful. (9) Departure from axisymmetry: Y—a departure from axisymmetry is clearly seen; N—no clear departure from axisymmetry is seen in the images. By departure from axisymmetry, I refer only to large-scale departures, and not to small-scale blobs, filaments etc. (10) My estimate of the compatibility of the PN departure from axisymmetry and the possibility that its binary companion caused this departure: (+)Compatible; (?)hard to tell. For example, for A 30 no departure is expected since the orbital period is very long, and indeed no large-scale departure from axisymmetry is observed, hence the + sign. For NGC 7008 a deviation from symmetry is expected and is indeed observed, hence a + sign. In K 1-14 a departure is observed, but the main signature is on the outskirts of the nebula, and hence an interaction with the ISM is possible. If the companion is associated with the PN central star, as claimed by Ciardullo *et al.*, then it can also cause a departure. Hence for this PN I put a question mark, since it is not clear if the morphology is compatible with the claimed companion, or it is solely due to an interaction with the ISM.

Four main processes can cause a large-scale deviation from axisymmetry. (i) Interaction with the ISM. In this case the most prominent features are on the outskirts on the nebula (e.g., Tweedy & Kwitter 1994, 1996), with smaller, or none at all, deviations from axisymmetry in the inner regions of the nebula. It is worth noting that in large PNs the ISM may penetrate the outer regions of the nebula, and influence the inner structure as well as the outer regions (Dgani & Soker 1998). (ii) A close companion in an eccentric orbit (Soker, Rappaport, & Harpaz 1998). This occurs when the companion is close enough to influence the mass loss process from the AGB star, and the eccentricity is large. (iii) A wide companion, with an orbital period in the range of $0.3\tau_{PN} \lesssim P_{\text{orb}} \lesssim 30\tau_{PN}$, where τ_{PN} is the formation time of the relevant part of the nebula. The formation time can be $\tau_{PN} \sim \text{several} \times 10^4$ yr for a halo, $\tau_{PN} \sim \text{several} \times 10^3$ yrs for the dense inner shell, and $\tau_{PN} \simeq \text{several} \times 100$ yrs for possible jets. For most of the PNs listed in Table 1, the relevant time is the expansion time of the nebula to its current radius $\tau_{PN} \sim R_{PN}/10 \text{ km s}^{-1} \sim 10^4 - 5 \times 10^4$ yrs. Another requirement of the latter mechanism is that

the velocity of the mass-losing star around the center of mass v_1 will be not too small. This means a companion of mass $M_2 \gtrsim 0.3M_\odot$, depending on the orbital separation (see §4). (iv) During the mass loss process itself, if one or a few long live cool (or hot) spots exist on the surface of the AGB star. This process seems to be important to massive stars (e.g., as suggested for the $\sim 30M_\odot$ star HD 179821 by Jura & Werner 1999), but it is not clear if it can operate efficiently in AGB stars, where the strong convection may not allow such spots to live long enough.

Simple estimates suggest that the deviation from axisymmetry, e.g., the dislocation of the central star from the center of the nebula, will be of the order of v_1/v_e , where v_e is the expansion velocity of the nebula. However, interaction of the wind with previously ejected mass from the AGB star may change the density of the wind in a different sense on both sides, and because of the sensitivity of emission to the density, a larger degree of departure from axisymmetry will be observed. The observed degree can be lower than expected if the direction of deviation is not perpendicular to the line of sight; if it is along the line of sight, the departure from axisymmetry will not be observed. In most cases I expect the degree of axisymmetry to be in the range from one to several times v_1/v_e .

In the next section I discuss the structure of each PN, and show that in most cases the PNs' structures can be used to strengthen the claims of Ciardullo *et al.* regarding the association, or non-association, of the putative companions with the central stars of the PNs.

3. NOTES ON THE INDIVIDUAL NEBULAE

In this section I analyze each PN separately. I will not discuss the position angle of the companion versus the direction of the departure from axisymmetry. This is because there are no details on the orbit (e.g., velocity), and the current position of the companion has little information about the orbit inclination, eccentricity, etc. In analyzing the different PNs, I use images from Balick (1987), Schwarz, Corradi, & Melnick, (1992), Acker *et al.* (1992), Manchado *et al.* (1996), and other sources as indicated below.

K1-14. There is a clear deviation from axisymmetry. The central star’s distance from the center of the nebula Δr is $\sim 15\%$ of the nebular radius, i.e., $\Delta r/R_{PN} \simeq 0.15$. For an expansion velocity of 15 km s^{-1} the age of the nebula is $\sim 2 \times 10^4$ yrs, just a little shorter than the orbital period. The velocity of the progenitor around the center of mass is relatively high, 0.5 km s^{-1} , and it can explain the departure from axisymmetry. Therefore, for this nebula the departure from axisymmetry may be attributed to the presence of the companion. However, this is not the only possibility. The central star is closer to the north-east part of the nebula, where a brighter arc is seen. This structure hints at an interaction with the ISM. Hence I put a question mark for this nebula (10th column in Table 1). Future observations will have to examine in more detail the properties of the companion and its association with the central star.

A 63. No clear deviation from axisymmetry is observed. This suggests that either the “companion” is not associated with the progenitor of the nebula (Ciardullo *et al.* marked it as a possible association), or that the real orbital separation is larger than the projected orbital separation, hence the orbital period is $\gtrsim 30$ times the age of the nebula. In any case, the absence of a clear departure from axisymmetry is compatible with the properties of the orbit even if the observed companion is a member of a binary system.

NGC 7008. A clear departure is observed, both in the outer and inner regions. This is expected, since the mass-losing star’s orbital velocity is large, 1.9 km s^{-1} , and the orbital period, although short, is within an order of magnitude of the formation time of this small nebula.

NGC 650-51. A clear asymmetry is seen along the waists of this bipolar PN. It cannot be attributed to an interaction with the ISM. The orbital velocity of $\sim 0.26 \text{ km s}^{-1}$ is too low to explain the large departure in the equatorial plane. The only way in which this “companion” could have caused the deviation is that it has a very eccentric orbit and currently it is close to its apastron position. In this case the orbital period is shorter, and when the companion was closer the orbital velocity of the mass-losing star was much higher. Although plausible, this is an unlikely explanation. More likely, a closer companion, but still at a wide orbital separation, exists, or a close companion which directly affects the mass-loss process orbits the central star with a

highly eccentric orbit (Soker *et al.* 1998). I marked this PN with a question mark.

PuWe 1. The images of this PN is from its discoverers, Purgathofer & Weinberger (1980), and Tweedy & Kwitter (1996). This PN shows a clear departure from symmetry, but with a bright arc on the outskirts of the nebula. This clearly suggests an interaction with the ISM, as was argued by Tweedy & Kwitter (1996), rather than a wide binary companion influence. This is compatible with the claim of Ciardullo *et al.* that the association of the observed “companion” with the central star is doubtful. Hence I marked it with a +.

NGC 2392. A small but clear departure is observed along the south-north direction, which is the long axis of the nebula, in the outer as well as the inner parts. Along the minor axis, the eastern side is longer by several percents than the western side. Such a deviation could not be caused by a companion at such a large orbital separation. However, the distance found by Ciardullo *et al.*, 6.41 kpc, is an upper limit. Most methods find a distance of $0.5 - 1.5$ kpc (Acker *et al.* 1992). Taking a distance to NGC 2392 to be ~ 1.2 kpc, the orbital period is $\sim 130,000$ yr, the orbital velocity is $v_1 \sim 0.5 \text{ km s}^{-2}$, and the companion will be less massive. These parameters for the companion and orbital motion can explain the observed departure from axisymmetry. Therefore, the morphology of this PN suggests that the companion is associated with the central star. Hence I marked it with a + sign, despite the classification of “possible association” by Ciardullo *et al.*. Another possibility is discussed together with NGC 1535 below.

NGC 1535. A small but clear asymmetry is observed along position angle $\sim 220^\circ$. Such a small departure is compatible with the parameters of the companion. Another possibility, for both this PN and NGC 2392, is that these are elliptical PNs with their long axis almost along the line of sight. The two sides are not identical due to an unequal collimated flows on the two sides of the long axis, which are not related to the presence of the companion. A study of the expansion velocities of these PNs along different directions is needed.

A 30. This PN has a very small departure from sphericity on its outskirts, which can safely be attributed to the ISM. No large-scale deviation is observed in the inner region, only blobs and filaments, hence I marked it to have no real deviation from axisymmetry. This is compatible with

the very long orbital motion and low velocity.

A 7. This a large nebula, which does not show a bright arc, is located far from the galactic plane and is surrounded by a tenuous ISM, hence it does not seem to interact with the ISM (Xilouris *et al.* 1996; Tweedy & Kwitter 1996). The distance to this PN is ~ 0.2 kpc, much smaller than the upper limit given by Ciardullo *et al.*, for which the orbital separation is ~ 200 AU and the orbital period $\sim 2,500$ yr. Such a binary should cause a noticeable departure. Although the nebula has an almost circular outer boundary, its inner region is elliptical, with unidentical bright regions on both sides. However, this is an old and large PN, and it is not clear whether this small deviation from axisymmetry is related to the presence of a companion, to instabilities which had time to grow, or to a weak interaction with the ISM. Therefore, I marked it with a question mark.

A 24. This PN has a point symmetric structure. The western side is a little more extended. I marked it as having a questionable large-scale deviation from point symmetric. Although the orbital period is long, this is a large PN, hence old one. Tweedy & Kwitter (1996) argue that it does not interact with the ISM. For all these, it is hard to tell whether its structure is compatible with the companion properties.

A 31. The image is taken from Tweedy & Kwitter (1994), who find that this PN interacts with the ISM. The morphology of this PN clearly comes from an interaction with the ISM. Any influence by the companion will be much smaller than the effect of the ISM. Therefore, I marked this PN with a question mark.

A33. A large deviation from axisymmetry, which cannot be attributed to the ISM since there is a deviation in the inner regions, and no bright arc is observed on the side closer to the central star. The distance of the star from the center of the nebula is $\Delta r/R_{PN} \simeq 0.07$. The departure of this PN is compatible with the properties of the companion, as already noted in a previous paper (Soker 1997).

NGC 6210. Ciardullo *et al.* find a distance smaller than 32 kpc. Most other methods give distances of ~ 2 kpc, i.e., a factor of 16 smaller. This means an orbital period $\lesssim 64$ times shorter,

and an orbital velocity of $\lesssim 0.8 \text{ km s}^{-1}$. Such a binary companion should cause a noticeable deviation from axisymmetry. This PN shows a small deviation from axisymmetry along its major axis. Since the deviation is not so clear, and Ciardullo *et al.* argue for a possible association between the companion and central star, I marked it with a question mark.

NGC 3132. This PN shows a clear asymmetry along position angle $\sim 45^\circ$. In that direction the nebula extends to a distance smaller by $\sim 20\%$ than the distance in the opposite direction. Such a large departure from axisymmetry is expected from a companion having the listed properties.

K 1-22. Although the image in Acker *et al.* (1992) is very faint, a small deviation from axisymmetry is observed in the inner region, more or less along the minor axis of the PN. This is compatible with the properties of the companion given in Table 1.

K 1-27. In Soker (1997) I suggested that the morphology indicates an interaction with the ISM. The finding of a companion by Ciardullo *et al.* suggests that it is the companion that causes the deviation from axisymmetry, or at least plays an important role in it. For this PN Ciardullo *et al.* argue that the companion is a WD. Hence I took its mass to be $0.6M_\odot$.

Mz 2. A clear asymmetry is observed: the two outer lobes are slightly displaced to the east relative to the inner bright shell, and a “missing” segment in the shell is observed at position angle $\sim 220^\circ$. This departure is compatible with the finding of Ciardullo *et al.*

Sp 3. In a previous paper (Soker 1997, Table 4) I noticed that the morphology of this PN suggests that it interacts with the ISM or contains a wide binary companion. The findings of Ciardullo *et al.* show that it is the companion that causes the departure from axisymmetry.

IC 4637. A departure from axisymmetry is observed, but it may come from an interaction with the ISM. I therefore marked it with a question mark.

4. PNs FOR WHOM NO COMPANIONS WERE FOUND

The purpose of the present paper is to demonstrate how the morphology of a PN can be used to strengthen (or not) the claim for the presence of a wide companion. However, in many cases a deviation from axisymmetry will be observed, but no companion will be found. These cases are:

(1) The deviation is caused by the motion of the PN through the ISM. I discussed this case in previous sections. Below a question arises in regard to A66 which has a clear asymmetry (Hua, Dopita & Mertinis 1998). However, this is a large low density nebula, and therefore the asymmetry is most probably due to the ISM.

(2) The projected separation of the companion and the PN central star is too small to be resolved, e.g., in the snapshot survey of Ciardullo *et al.*, the resolution with the HST is 0.1 arcsec. This will happen if the orbital separation is small, or if the orbital separation is large but the binary plane is almost edge-on, and the companion line of sight happens to be closed to that of the central star. Note that to observe a deviation from axisymmetry the orbit must not be face on, so in many cases the orbit will be close to being edge-on.

(3) The companion is a low mass main sequence star, and it is too faint to be detected. In Table 1, 18 out of 19 companions, or possible companions, have a mass of $\gtrsim 0.5M_{\odot}$. Therefore, we can safely deduce that the study of Ciardullo *et al.* is limited mainly to main sequence companions of masses $\gtrsim 0.5M_{\odot}$ (we return to this point below).

(4) The deviation from axisymmetry is due to a close companion at an eccentric orbit (Soker *et al.* 1998). Such systems are likely to form Bipolar PNs.

(5) A companion that enters the envelope of the progenitor of the PN may cause a deviation from axisymmetry as well. This has been suggested as a possible explanation for the displacement of the two outer rings of SN1987A (Soker 1999). Regarding mechanisms (4) and (5), a clear deviation from axisymmetry is seen in the bipolar PN NGC 2346, for which there is a very close companion with an orbital period of 16 days (Bond & Livio 1990).

(6) In the survey of Ciardullo *et al.* PNs located in a high field star density were also classified as non-detection, since a wide binary companion could not be statistically identified in these cases.

It is possible therefore, that the companions of some of the non-detection PNs which have nebular departure from axisymmetry are seen in the survey, but can not be statistically distinguished from the field stars.

With these possibilities in mind, I examine the PNs from the snapshot survey of Ciardullo *et al.* for which no companions have been detected. Here again, I consider a PN to have a clear departure from axisymmetry when the scale of departure is larger than the typical size of blobs and filaments in the PN. The list of 67 PNs and their properties was kindly supplied to me by R. Ciardullo. In this list there are 4 PNs which are classified as Bipolar, and are likely to be formed from a close binary interaction (Soker 1997, Table 3). Considering point (4) above, I do not consider these PNs. Out of these 67 PNs, 13 PNs were claimed by Soker (1997, Table 4) to be descendants of common envelope evolution. Following point (5) above, I do not consider these PNs.

Out of the rest 50 PNs, The deviations from axisymmetry of 15 PNs were noted by Soker (1997) as follows. For 9 PNs the deviation from axisymmetry seems to result from interaction with the ISM; two out of these 9 show no deviation in the inner regions, and therefore have no influence of a wide companion (A16; NGC 6894), while for the other 7 it is hard to tell (due to, e.g., an extended filamentary structure). For NGC 5979 Soker (1997) left it open whether the departure from axisymmetry is due to an interaction with the ISM or to a wide companion. IC 4593 show both signs of interaction with the ISM and a wide companion. For 4 PNs Soker (1997) attributes the departure to a wide companion (IC 5148-50; NGC 2022; NGC 3242; NGC 7662). In these PNs the departure from axisymmetry is most prominent in the inner regions, and therefore can be caused by a companion with a relatively short orbital period. Soker, Zucker, & Balcik (1992) attributed the departure from axisymmetry of NGC 3242 to a star with an orbital period of $\sim 4,000$ yrs, but a shorter period is also possible. A shorter orbital period is possible for NGC 2022, for which the deviation is of the inner “jets”. A careful examination of images in the literature reveals 3 other PNs with a clear departure from axisymmetry (NGC 6153, Gorny *et al.* 1999; NGC 5882, NGC 6804 Schwarz *et al.* 1992), and 2 with departure which I cannot tell if

compatible with a wide companion (NGC 6891, Chu, Jacoby & Arendt 1987; NGC 2792, Schwarz *et al.* 1992).

For 23 PNs out of the 50 I consider, I could not tell whether they have a departure from axisymmetry which is compatible with a wide companion (e.g., they have interaction with the ISM; they have filamentary structure with small number of large filaments; no high resolution image exists). Out of the rest 27 PNs, 8 show clear departure, and for 3 it is not clear (these 11 PNs were discussed in the previous paragraph). I conclude that $\sim 35 \pm 5\%$ of PNs for whom no companions were found by Ciardullo *et al.*, show departure from axisymmetry compatible with the presence of a wide stellar companion. This should be compared to the 10 PNs with probable detection, from which 8 have departure compatible with the claimed companion (both ‘Y’ and ‘+’ in Table 1). Considering the detection limits, and in particular the companion mass of $\sim 0.5M_{\odot}$, I find the differences in detection, 80% compared with 35%, significant, despite the small number of PNs.

I return now to the question of the limit on the companions masses. I used the I-band limiting magnitude (supplied by Ciardullo) of the 8 PNs I claim to have wide companions and the 3 PNs with question marks, to estimate the upper limit on the masses of possible main sequence companions. I took distances from Cahn, Kaler & Stanghellini (1992), and the mass-luminosity relation for main sequence stars. Because of the strong dependence of luminosity on mass in that relation, the calculation is quite robust to the uncertainties in the different quantities entering the calculation. The uncertainties in the distances may be up to a factor of ~ 3 , which introduce uncertainties of up to $\sim 70\%$ in the limiting masses. Because of these uncertainties I do not find it appropriate to list each object separately (unlike the case of the PNs presented in Table 1, for which Ciardullo *et al.* used the companions to determine the distances). Overall, I find the limiting masses to be around $\sim 0.5M_{\odot}$ for most objects. This is compatible with the masses of detected companions (Table 1).

A detailed statistical study of binary systems that cause departure from axisymmetry using a population synthesis code, is planned for a forthcoming paper. The code that will be used is

the one used by Rappaport & Soker (1999), from which I take the different values used below. I crudely estimate the number of expected wide binaries as follows. For an orbital separation smaller than ~ 1 AU (an orbital period of one year), a strong interaction of the binary system will form a bipolar PNs, or the system will enter a common envelope. I exclude such systems here. For the rest, the distribution of binary orbital period is uniform in $\log(P)$, up to $P \simeq 10^6$ yrs. The orbital velocity of the mass-losing star, of mass M_1 , around the center of mass with a companion of mass M_2 , is (assuming a circular orbit)

$$v_1 = 0.4 \left(\frac{P}{10^4 \text{ yr}} \right)^{-1/3} \left(\frac{M_1 + M_2}{1 M_\odot} \right)^{-2/3} \left(\frac{M_2}{0.3 M_\odot} \right) \text{ km s}^{-1}. \quad (1)$$

As mentioned before, an orbital velocity of $\sim 0.3 \text{ km s}^{-1}$ may cause a noticeable departure from axisymmetry. The mass distribution in binary systems is somewhat peaked toward $M_2/M_1 = 1$, hence most companions of PN progenitors will have masses of $M_2 \gtrsim 0.3 M_\odot$. This means that binaries with orbital periods of up to $\text{few} \times 10^5$ yrs can be counted, especially if eccentric orbits are considered as well. From below, we require that the orbital period will not be too short, this gives an orbital period longer than ~ 500 yrs. Overall, most companions in a logarithmic interval of ~ 2 , which is $\sim 1/3$ of all binary systems may cause deviation from axisymmetry. The question is what is the fraction of binary systems among all progenitors of PNs. This fraction is taken by different authors to be in the range of 0.6-1 (Rappaport & Soker 1999). I conclude that wide binaries will cause a clear departure from axisymmetry in 20 – 35% of all PNs. The fraction of PNs below the detection limit mentioned above, I crudely estimate to be half this number 10 – 17%. Adding to these some interaction with the ISM (e.g., a dense cloud), and large scale instabilities in the mass loss process, which I interpreted as wide binaries in analyzing the PNs in this section, I find the fraction of PNs with departure from pure axisymmetry among the non-detection PNs ($\sim 35\%$) to be quite reasonable.

5. SUMMARY

Stellar companions to progenitors of PNs will influence the morphology of the descendant PNs. Close companions will affect directly the mass loss process from the AGB progenitor. Such close binaries are not likely to be resolved. Very wide binaries (as well as the wide binaries discussed here) may form a small bubble inside the nebula (Soker 1996). In the intermediate range, binary systems which have orbital periods in the range of several $\times 100$ to a few $\times 10^5$ years, may cause the PN to have a large-scale departure from axisymmetrical structure. The goal of the present paper was to show that the structure of a PN can be used to strengthen an argument for an association, or a non-association, of a putative wide companion with the progenitor of the PN. In a recent work Ciardullo *et al.* (1999) list 19 PNs for which they argue a probable, possible, or doubtful association of stars with the progenitors of the PNs. Ciardullo *et al.* did not use the structure of the PNs as a tool to further support their claims. In the present paper I used their list, and demonstrated that the structures of the PNs are compatible in most cases with their claims, hence this paper strengthens them. The greatest confusion may come from a possible influence of the ISM on the PN morphology, especially for large PNs. I tried to present arguments for and against ISM interaction where possible.

In the opposite direction, PNs whose central stars have companions can be used to further study the influence of wide companions on the PNs morphology. For this, a determination of the binary component velocity is needed. Very helpful but more difficult to determine will be the eccentricity and inclination of the orbital plane.

From a list of 27 PNs observed with HST for which no companions were found by Ciardullo *et al.*, I claimed that 8 PNs are likely to have companions which have caused their structures to depart from axisymmetry. In addition to 5 PNs from this list which were predicted to have wide binary companions by Soker (1997), I added here to the list three PNs: NGC 5882, NGC 6153, and NGC 6804. I estimated that the observations of Ciardullo *et al.* will not detect main sequence companions of masses below $\sim 0.5M_{\odot}$ in these 8 PNs. A search for a wide binary companion in each of these 8 PNs is encouraged. For three PNs out of the 27 I could not tell if a companion is

compatible or not with their structure. That only 8 of the 27 PNs have morphology compatible with a wide binary companion, compared with 8 out of 10 PNs that were claimed to have probable companion by Ciardullo *et al.*, strengthen the arguments presented in the present paper.

ACKNOWLEDGMENTS: I would like to thank the referee, Robin Ciardullo, for his suggestion to study the sample of PNs for whom no companions were found, and for supplying the data for these PNs, and Matthew Bobrowsky for a careful reading of the manuscript. This research was supported in part by a grant from the University of Haifa and a grant from the Israel Science Foundation.

REFERENCES

- Acker, A., Ochsenbein, F., Stenholm, B., Tyllenda, R., Marcout, J., & Schohn, C. 1992, Strasbourg-ESO Catalogue of Galactic Planetary Nebulae, (Pub. by ESO)
- Balick, B. 1987, *AJ*, 94, 671.
- Bobrowsky, M., Sahu, K. C., Parthasarathy, M., & Garcia-Lario, P. 1998, *Nature*, 392, 469.
- Bond, H., & Livio, M. 1990, *ApJ*, 355, 568.
- Cahn, J. H., Kaler, J. B., & Stanghellini, L. 1992, *A&AS*, 94, 399.
- Chu, Y.-H., Jacoby, G. H., & Arendt, R. 1987, *ApJSupp.*, 64, 529.
- Ciardullo, R., Bond, H. E., Sipior, M. S., Fullton, L. K., Zhang, C.-Y., & Schaefer, K. G. 1999, *AJ*, in press (astro-ph/9904043).
- Corradi, R. L. M., Perinotto, M., Villaver, E., Mampaso, A., & Goncalves, D. R. 1999, *ApJ*, in press.
- Dgani, R., & Soker, N. 1998, *ApJ*, 495, 337.
- Gorny, S. K., Schwarz, H. E., Corradi, R. L. M., & Van Winckel, H. 1999, *A&AS*, 136, 145.
- Hua, C. T., Dopita, M. A., & Martinis, J. 1998, *A&AS* 133, 361.
- Jura, M., & Werner, M. W. 1999, preprint.
- Manchado, A., Guerrero, M., Stanghellini, L., & Serra-Ricart, M. 1996, The IAC Morphological Catalog of Northern Galactic Planetary Nebulae (Tenerife: IAC).
- Purgathofer, A., & Weinberger, R. 1980, *A&A*, 87, L5.
- Rappaport, S., & Soker, N. 1999, preprint.
- Schwarz, H. E., Corradi, R. L. M., & Melnick, J. 1992, *A&A, Supp. Ser.* 96, 23.
- Soker, N. 1994, *MNRAS*, 270, 774.
- Soker, N. 1996, *MNRAS*, 283, 1405.
- Soker, N. 1997, *ApJSupp.*, 112, 487

- Soker, N. 1999, MNRAS, 303, 611.
- Soker, N., Rappaport, S., & Harpaz, A. 1998, ApJ, 496, 842.
- Soker, N., Zucker, D. B., & Balick, B. 1992, AJ, 104, 2151.
- Tweedy, R. W., & Kwitter, K. B. 1994, AJ, 108, 188.
- Tweedy, R. W., & Kwitter, K. B. 1996, ApJ Supp., 107, 255.
- Xilouris, K. M., Papamastorakis, J., Paleologou, E., & Terzian, Y. 1996, A&A, 310, 603.

TABLE 1
COMPANIONS AND ORBITAL MOTION

Object	PN G	$M_2(M_\odot)$	$a_{\text{pr}}(\text{AU})$	$P_{\text{orb}}(\text{yr})$	$v_1(\text{km s}^{-1})$	$D_{PN}(\text{pc})$	B	D	C
K 1-14	045.6+24.3	0.7	1080	29300	0.5	0.68	P	Y	?
A 63	053.8-03.0	1.5	3440	133000	0.5	0.24	S	N	+
NGC 7008	093.4+05.4	1.1	160	1470	1.9	0.15	P	Y	+
NGC 650-51	130.9-10.5	0.8	5520	323000	0.3	1.34	D	Y	?
PuWe 1	158.9+17.8	0.5	1620	56500	0.3	1.82	D	Y	+
NGC 2392	197.8+17.3	1.7	<16970	<1400000	>0.2	<0.61	S	Y	+
NGC 1535	206.4-40.5	0.8	2400	93500	0.4	0.23	P	Y	+
A 30	208.5+33.2	0.8	10580	859000	0.2	1.24	S	N	+
A 7	215.5-30.8	0.6	<11520	<1060000	>0.1	<46.8	S	Y	?
A 24	217.1+14.7	0.5	7920	607000	0.2	4.1	D	Y?	?
A 31	219.1+31.2	0.05	<115	<1340	>0.2	<2.09	P	Y	?
A 33	238.0+34.8	0.5	2110	84200	0.3	1.52	P	Y	+
NGC 2610	239.6+13.9	1.0	<19630	<2070000	>0.2	<5.93	S	Y?	?
NGC 3132	272.1+12.3	3.1	1310	24100	1.3	0.11	P	Y	+
K 1-22	283.6+25.3	0.6	470	8510	0.7	1.16	P	Y	+
K 1-27	286.8-29.5	0.6	260	3540	0.9	0.10	P	Y	+
Mz 2	329.3-02.8	1.9	600	9020	1.4	0.24	P	Y	+
Sp 3	342.5-14.3	1.1	740	14500	0.9	0.41	P	Y	+
IC 4637	345.4+00.1	1.6	1210	26900	0.9	0.05	S	Y	?